arXiv:1208.1694v1 [q-bio.PE] 5 Aug 2012

Stochastic dynamics of dengue epidemics

Tânia Tomé[1], Suani T. R. Pinho[2], Florisneide R.
Barreto[3], David R. Souza[1], and Mário J. de Oliveira[1]
[1]*Instituto de Física, Universidade de São Paulo,*
Caixa Postal 66318, 05314-970, São Paulo, Brazil
[2]*Instituto de Física, Universidade Federal da Bahia, 40210-340, Salvador, Brazil*
[3]*Instituto de Saúde Coletiva, Universidade Federal da Bahia, 40110-140, Salvador, Brazil*

Abstract

The spreading of vector-transmitted diseases, like dengue, is a relevant problem of public health. Different techniques are used to understand its complex dynamics. In this paper we use a stochastic Markovian dynamics approach to describe the spreading of dengue and the threshold of the disease. The coexistence space is composed by two structures representing the human and mosquito populations. The human population follows a susceptible-infected-recovered (SIR) type dynamics and the mosquito population follows a susceptible-infected-susceptible (SIS) type dynamics. The human infection is caused by infected mosquitoes and vice-versa so that the SIS and SIR dynamics are interconnected. After setting up the master equation associated to the stochastic dynamics we develop a truncation scheme to solve the evolution equations from which we get the epidemic curve and the threshold of the disease as well as the reproductive ratio. The results are compared to the actual outbreaks of the disease.

PACS numbers: 87.10.Mn, 87.10.Hk, 05.70.Fh, 05.70.Ln

I. INTRODUCTION

Dengue is a vector-borne infectious disease with very complex dynamics, whose spreading is a relevant problem of public health. The disease is transmitted to human mainly by the mosquito *Aedes Aegypti*. Many factors are determinant for the transmission of dengue in urban centers such as the climatic conditions for the vector proliferation, the human concentration and mobility. Although many efforts are put in the development of a vaccine against the four types of virus, until now the only available strategy to reduce the spreading of the disease [1] is the control of the vector population. Therefore, it is very important to analyze the effect of vector control avoiding the occurrence of dengue epidemics. In this context, the inter-host modeling of dengue dynamics and control may be a very useful tool for helping its understanding and the establishment of vector control strategies.

Different techniques and approaches are used to model the dynamics of transmitted diseases [2, 3] such as deterministic differential equations [4], stochastic dynamics [5–7], cellular automata [8], and complex networks [9]. Concerning vector transmitted diseases including dengue modeling, there are also different schemes and approaches [2, 3] based on deterministic models of differential equations [10–12], probabilistic cellular automata [13–16] and complex networks [17, 18]. However, as far as we know, it is the first time that a description in terms of a master equation defined on a lattice is used on the investigation of a vector-borne infectious disease although it was used by some of us to investigate epidemic models of direct transmitted diseases [5].

As in other works [10–12, 15], the present approach is also motivated by actual data of dengue epidemics [19] in particular, by two outbreaks of dengue occurred in Salvador, Bahia, Brazil in 1995 (without vector control) and 2002 (with vector control). Those data had also motivated a previous analysis based on the basic reproductive ratio [10].

The first dengue model was proposed by Newton and Reiter [20] in 1992 assuming a susceptible, exposed, infected and recovered (SEIR) structure for humans and susceptible, exposed and infected (SEI) structure for mosquitoes due to the fact that the mosquitoes die before being removed. This framework has been followed by other continuous and discrete dengue models [10–12, 14, 15]. Here we consider a simpler model, illustrated in figure 1, assuming a susceptible, infected and recovered (SIR) structure for humans and susceptible, infected and susceptible (SIS) structure for mosquitoes. The infection of humans is due to



FIG. 1: Illustration of the reactions. Left panel: SIR structure. A susceptible human (H_S) becomes infected (H_I) through a catalytic reaction mediated by infected mosquitoes (M_I). An infected human becomes recovered (H_R) spontaneously and remains permanently in this state. Right panel: SIS structure. A susceptible mosquito (M_S) becomes infected through a catalytic reaction mediated by infected humans. An infected mosquito spontaneously becomes susceptible.

mosquitoes and the infection of mosquitoes is due to humans. In other words the infection reactions, S→I, on both structures are catalytic and not autocatalytic as happens to the original SIR and SIS models [5–7]. The other reactions, I→R in the SIR structure and I→S in the SIS structure, are spontaneous reactions. The mosquito structure has been simplified by suppressing the death and the birth of mosquitoes which amounts to saying that a dead mosquito is immediately replaced by a new-born (susceptible) mosquito. We are, therefore, assuming that the number of mosquitoes remains constant throughout the outbreak of the epidemics.

The features of our model that differ from that of Newton and Reiter [20] are as follows. Firstly, we do not distinguish between susceptible and exposed states both for humans and mosquitoes. Secondly, deaths of humans are not considered since the human life time is much larger than the period of the disease. As to the deaths of the mosquitoes, they are implicit in the model in the following sense. The reaction $I \to S$ for the mosquitoes is to be interpreted as the death of an infected mosquito and the simultaneous birth of a susceptible mosquito. The major difference however rests on the use of a stochastic lattice model, which takes into account the spatial distribution of humans and mosquitoes.

This paper is organized as follows. In section II we introduce the model and derive from the master equation the evolution equation for the densities. We also define the quantities that characterize the spreading of the epidemics. In section III we develop the simple mean-field analysis. In section IV we introduce the evolution equation for the two-site probabilities and perform a pair mean-field analysis. Also in this section we compare the results coming from the pair mean-field with actual data coming from the outbreak of the dengue disease occurred in Salvador in 1995 and 2002. Finally, concluding remarks are presented in the last section.

II. MODEL AND EVOLUTION EQUATIONS

The modeling of disease spreading that we consider here corresponds to a continuous time stochastic Markovian process defined on a lattice where the sites are occupied by human individuals or by mosquitoes. In order to properly describe the human and mosquito populations we consider two sublattices of the whole lattice, one for each population. The sublattices, named H and M, are interpenetrating in such a way that the nearest neighbor sites of a site of one sublattice belong to the other sublattice. Each site of the sublattice M can be in one of two states, either occupied by a susceptible mosquito M_S or by an infected mosquito M_I. Each site of the sublattice H can be in one of three states, occupied by a susceptible human H_S, occupied by an infected human H_I or occupied by a recovered human H_R. The system evolves in time according to the following stochastic dynamics.

At each time step a site of the lattice is chosen at random. If the site belongs to sublattice H then the following update is used. (a) If the chosen site is occupied by a susceptible human then it becomes infected with rate a times the fraction of infected mosquitoes in its neighborhood. If the chosen site is occupied by an infected human then it becomes recovered spontaneously with rate c. Once recovered the individual remains permanently in this state which means that, if the chosen site is occupied by a recovered human, it remains unchanged. (b) Next, we consider the case in which the chosen site belongs to the M sublattice. If it is occupied by a susceptible mosquito then it becomes infected with a rate b times the fraction of infected humans in its neighborhood. If the chosen site is occupied by an infected mosquito it spontaneously becomes susceptible with rate e. Our simple model is therefore described by four parameters a, b, c and e. In the applications we will further simplify by setting $b = a$.

For convenience we introduce a stochastic variable η_i associated to each site i of the lattice that takes the values 0, 1, 2, 3, or 4 according to whether the site i is occupied, respectively, by a susceptible mosquito (M_S), an infected mosquito (M_S), a susceptible human (H_S), an infected human (H_I) or a recovered human (H_R). The time evolution of the probability distribution $P(\eta)$ of configuration $\eta = \{\eta_i\}$ is governed by the master equation

$$\frac{d}{dt}P(\eta) = \sum_i \{w_i(\mathcal{A}_i^- \eta)P(\mathcal{A}_i^- \eta) - w_i(\eta)P(\eta)\}. \tag{1}$$

where $w_i(\eta)$ is the transition rate from η to $\eta' = \mathcal{A}_i\eta$ and $\mathcal{A}_i$ is the operator that changes $\eta_i \to \eta_i'$ as follows: 0→1, 1→0, 2→3, 3→4; and $\mathcal{A}_i^-$ is the inverse of $\mathcal{A}_i$. The transition rate $w_i(\eta)$ is defined according to the rules stated above. The time evolution of an average $\langle f(\eta)\rangle = \sum_\eta f(\eta)P(\eta)$ is obtained from the master equation and is given by

$$\frac{d}{dt}\langle f(\eta)\rangle = \sum_i \langle [f(\mathcal{A}_i\eta) - f(\eta)]w_i(\eta)\rangle. \tag{2}$$

Instead of using the full probability distribution $P(\eta)$ we consider an equivalent description in terms of the various marginal probability distribution, obtained from $P(\eta)$. The time evolution of the several marginal probability distributions are obtained from the master equation (1) and comprises a hierarchic set of coupled equations which is equivalent to the master equation. This approach is convenient because it allows to obtain a solution of the set of equations by a truncation scheme to be explained shortly. In what follows we assume translation invariance and isotropy.

Let us denote by the $P(\eta_i)$ the marginal one-site probability that represents the probability that a site i is in state η_i and by $P(\eta_i, \eta_j)$ the marginal two-site probability that represents the probability that site i is in state η_i and a neighboring site j is in state η_j. Other marginal probabilities are denoted in an analogous way. The evolution equation of the one-site probability $P(1)$, which represents the density of infected mosquitoes, can be obtained from equation (2) if we recall that $P(1) = \langle \delta(\eta_i, 1)\rangle$ and is given by

$$\frac{d}{dt}P(1) = bP(03) - eP(1). \tag{3}$$

where we used the relation $P(03) = \langle \delta(\eta_i, 0)\delta(\eta_j, 3)\rangle$. The equation for $P(0)$, the density of susceptible mosquitoes, can be obtained from equation (3) by using the property

$$P(0) + P(1) = 1. \tag{4}$$

The evolution equations for $P(2)$ and $P(3)$, the densities of susceptible and infected humans, respectively, are obtained similarly from equation (2) and are given by

$$\frac{d}{dt}P(2) = -aP(12), \tag{5}$$

$$\frac{d}{dt}P(3) = aP(12) - cP(3). \tag{6}$$

The equation for $P(4)$, the density of recovered humans, can be obtained from equations (5) and (6) by taking into account the property

$$P(2) + P(3) + P(4) = 1. \tag{7}$$

To characterize the threshold of the epidemic it is convenient to write equations (3) for the density of infected mosquitoes and (6) for the density of infected individuals in the form

$$\frac{d}{dt}P(1) = bP(0|3)P(3) - eP(1), \tag{8}$$

$$\frac{d}{dt}P(3) = aP(2|1)P(1) - cP(3), \tag{9}$$

where $P(2|1) = P(12)/P(1)$ is the conditional probability of occurrence of a susceptible individual given an infected neighboring mosquito and $P(0|3) = P(03)/P(3)$ is the conditional probability of occurrence of a susceptible mosquitoes given an infected neighboring individual. Using the simplified notation $x = P(1)$ and $z = P(3)$ the set of equations (8) and (9) can be written as

$$\begin{pmatrix} dx/dt \\ dz/dt \end{pmatrix} = \begin{pmatrix} -e & bP(0|3) \\ aP(2|1) & -c \end{pmatrix} \begin{pmatrix} x \\ z \end{pmatrix}, \tag{10}$$

where $P(0|3)$ and $P(2|1)$ are the cross transmission probabilities. At the early stages of the epidemic the cross transmission probabilities $P(0|3)$ and $P(2|1)$ can be considered to be constant (independent of time) and the set of equations (8) and (9) becomes a linear set of equations. A fundamental quantity that characterizes the spreading of the disease is the so-called reproductive ratio R_0 which is defined here as follows

$$R_0 = \frac{ab}{ce}P(2|1)P(0|3). \tag{11}$$

The threshold of epidemic is determined by the largest eigenvalue of the matrix (10), which is related to the reproductive ratio by

$$R_0 = \left(1 + \frac{\lambda}{e}\right)\left(1 + \frac{\lambda}{c}\right), \tag{12}$$

or by

$$\lambda = \frac{1}{2}\{-(e + c) + \sqrt{(e - c)^2 + 4ecR_0}\}. \tag{13}$$

According to the linear analysis the threshold of epidemic occurs when the largest eigenvalue vanishes which happens, according to equation (12), when $R_0 = 1$. Moreover, when $\lambda < 0$ there is no transmission of disease. According to equation (12) this happens when $R_0 < 1$. The spreading of the disease occurs when $\lambda > 0$, that is, when $R_0 > 1$. The reproductive ratio characterizes not only the threshold of the disease but also its strength.

III. SIMPLE MEAN-FIELD APPROXIMATION

The three equations for the time evolution of $P(1)$, $P(2)$ and $P(3)$ are not a set of closed equations because they depend on the two-site probabilities $P(12)$ and $P(03)$. To get closed equations we firstly use a scheme, called simple mean-field (SMF) approximation, which consists in writing a two-site probability as the product of one-site probabilities, that is, $P(\eta_i, \eta_j) = P(\eta_i)P(\eta_j)$. Using the abbreviations $P(1) = x$, which is the density of infected mosquitoes, $P(2) = y$, which is the density of susceptible individuals, and $P(3) = z$, which is the infected individuals. This approximation gives $P(12) = P(1)P(2) = xy$ and $P(03) = P(0)P(3) = [1-P(1)]P(3) = (1-x)z$ so that equations (3), (5) and (6) are reduced to the following forms

$$\frac{dx}{dt} = b(1-x)z - ex, \tag{14}$$

$$\frac{dy}{dt} = -axy, \tag{15}$$

$$\frac{dz}{dt} = axy - cz. \tag{16}$$

To describe the spreading of the disease we consider an initial condition such that all individuals are susceptible so that $y = 1$ and $z = 0$. Moreover we consider a very small density of infected mosquitoes, that is, $x = \epsilon$ where $\epsilon << 1$. Then we look for solutions for x, y and z in the limit $t \to \infty$. In this limit the density of infected mosquitoes and infected individuals vanish, $x \to 0$ and $z \to 0$. If the density of recovered individuals $\rho = 1 - y - z$ approaches a nonzero value than the disease spreads. On the other hand ρ approaches zero (when $\epsilon \to 0$) the disease does not spread. This means that the stationary solution $x = 0$, $z = 0$, $y = 1$ ($\rho = 0$) is the solution corresponding to the non-spreading regime. To obtain the stability of this solution we linearize the above equations around this solution to obtain

$$\frac{dx}{dt} = bz - ex, \tag{17}$$

$$\frac{d\rho}{dt} = ax, \tag{18}$$

$$\frac{dz}{dt} = ax - cz. \tag{19}$$

A linear analysis of stability amounts to determine the eigenvalues of the matrix composed by the coefficients of the right hand side of these equations, given by

$$\begin{pmatrix} -e & 0 & b \\ a & 0 & 0 \\ a & 0 & -c \end{pmatrix}. \tag{20}$$

One eigenvalue is zero and the others are the roots of

$$\lambda^2 + (e+c)\lambda + ec - ab = 0, \tag{21}$$

that is,

$$\lambda_\pm = \frac{1}{2}\{-(e+c) \pm \sqrt{(e-c)^2 + 4ab}\}. \tag{22}$$

The solution is stable as long as $\lambda_+ < 0$, that is when $ec > ab$. The threshold of the spreading occurs when

$$\frac{ab}{ec} = 1. \tag{23}$$

For $a = b$, the threshold line is described by

$$\frac{a^2}{ec} = 1, \tag{24}$$

as shown in the phase diagram e/c versus a/c of figure 2a.

The threshold obtained by the above linear analysis can equivalently be obtained from the condition $R_0 = 1$, that is, when the reproductive ratio R_0 given by equation (11) equals the unit. In the present case of simple mean-field approach $P(2|1) = P(2)$ and $P(0|3) = P(0)$. The initial condition gives, $P(2) = 1$ and $P(0) = 1$ so that

$$R_0 = \frac{ab}{ce}, \tag{25}$$

which coincides with the condition (23) when $R_0 = 1$. It is worth mentioning that R_0 depends on the parameters a, b, c and e only through the ratios e/c, b/c and a/c. Figure 2b shows the lines at constant R_0 in the plane e/c versus a/c for the case $b = a$ as well as the threshold of epidemic characterized by $R_0 = 1$.



FIG. 2: Phase diagram in the plane e/c versus a/c showing the regions of spreading (S) and non-spreading (NS) of the disease for the (a) SMF approximation and (b) PMF approximation. The continuous curve is the threshold of spreading of disease. When $e \to 0$, the value of a/c approaches zero for the SMF and the value $a/c = 0.5$ for the PMF. The curves inside the S region are lines of constant reproductive ration R_0, from top to bottom: R_0 =1.2, 1.5, 2, 4, and 10. The threshold of spreading of disease (continuous line) corresponds to $R_0 = 1$.

To obtain the time dependent properties, including the epidemic curve we have solved the set of equations numerically starting from the initial condition $x = \epsilon$, $y = 1$ and $z = 0$, where $\epsilon = 10^{-6}$. An epidemic is characterized by the epidemic curve defined as the number of cases occurring in unit time, for instance, in a day or in a weak, plotted as a function of time. In other terms it is the number of susceptible individuals that are being infected per unit time. Instead of number of individual we may define the epidemic curve as the density of individuals that are being infected per unit time $\zeta = -dy/dt$. Figure 3a shows an example of the epidemic curve obtained from the SMF approximation together with the density of infected mosquitoes.

For sufficiently large values of time the density of recovered individuals ρ approaches its maximum value. Notice that the final density of recovered individuals ρ depends on a, c and e only through the ratios a/c and e/c and is the integral of the epidemic curve. Indeed, if we integrate $\zeta = -dy/dt$, we get

$$\int_0^\infty \zeta dt = \rho, \tag{26}$$

where we used the initial condition $y(0) = 1$ and the result that the final density of infected individuals vanish, that is, $z(\infty) = 0$, and the constraint $\rho + y + z = 1$. The final density of recovered individual, which is a measure of the size of the epidemics, may be understood in this approach as the order parameter in the sense that it vanishes in the non-spreading regime and is nonzero in the regime where the disease spreads.

Near the threshold of epidemics the density of infected mosquitoes is much smaller than unit and we may neglect x in the first term on the right hand side of equation (14). The evolution equation for x becomes then

$$\frac{dx}{dt} = bz - ex, \tag{27}$$

which together with equations (15) and (16) allows us to determine explicitly the size of the epidemics ρ. This is possible because these equations implies a conservation law obtained as follows. We start by defining the quantity $\phi = cx + bz$. From equations (27) and (16) its evolution equation is given by

$$\frac{d\phi}{dt} = -cex + abxy. \tag{28}$$

The ratio of equations (28) and (15) gives

$$\frac{d\phi}{dy} = \frac{ce}{ay} - b \tag{29}$$

which can be integrated to give

$$cx + bz = \frac{ce}{a} \ln y + b(1 - y) \tag{30}$$

which is the desired conservation law. The constant of integration was obtained by remembering that at $t = 0$, $x = \epsilon \to 0$, $y = 1$ and $z = 0$.

When $t \to \infty$, $x = 0$, $z = 0$ and $1 - y = \rho$ so that

$$\ln(1 - \rho) + R_0 \rho = 0 \tag{31}$$

where we used the relation $R_0 = ab/ce$. This equation determines the size of the epidemics ρ and may be written in the form

$$1 - \rho = e^{-R_0 \rho} \tag{32}$$

This equation is the same equation obtained by Kendal [21] for the model introduced by Kermack and McKendrick [22] to describe the directed transmitted epidemics [2, 3] and also obtained by means of the simple mean-field approach to the SIR model on a lattice [7]. Near the threshold it is given by $\rho = 2(R_0 - 1)$.

IV. PAIR MEAN-FIELD APPROXIMATION

A. Evolution equations

Next we set up equations for the two site probabilities. The evolution equation for the probability $P(03)$ of a site being occupied by a susceptible mosquito and a neighboring site by an infected individual is given by

$$\frac{d}{dt}P(03) = eP(13) - (c + rb)P(03) + (1 - r)aP(021) + (1 - r)bP(303), \tag{33}$$

where $r = 1/k$, and k is the coordination number of the lattice. The notation $P(\eta_i, \eta_j, \eta_k)$ stands for the joint three-site probability where i and k are nearest neighbor sites of j. The evolution equation for the probability $P(12)$ of a site being occupied by an infected mosquito and a neighboring site by a susceptible individual is given by

$$\frac{d}{dt}P(12) = -(e + ra)P(12) + (1 - r)bP(302) + (1 - r)aP(121). \tag{34}$$

The equations for the other two-site probabilities, $P(02)$, $P(04)$, $P(13)$, and $P(14)$ are not necessary because of the following relations

$$P(02) + P(03) + P(04) = P(0), \tag{35}$$

$$P(12) + P(13) + P(14) = P(1), \tag{36}$$

$$P(02) + P(12) = P(2), \tag{37}$$

$$P(03) + P(13) = P(3), \tag{38}$$

$$P(04) + P(14) = P(4). \tag{39}$$

Let us consider now the equations for the time evolution of $P(1)$, $P(2)$, $P(3)$, given by equations (3), (5) and (6), and the equations for time evolution equations of $P(03)$ and $P(12)$, given by equations (33) and (34). These five equations are not closed equations because (33) and (34) depend on three-site probabilities. To get a set of closed equations we use now a truncation scheme at the level of two-site probabilities [23]. To this end we start by writing the three-site probability as $P(\eta_i, \eta_j, \eta_k) = P(\eta_i, \eta_k|\eta_j)P(\eta_j)$ where i and k are distinct neighbor sites of site j. Next we use the following approximation for the conditional probability $P(\eta_i, \eta_k|\eta_j) = P(\eta_i|\eta_j)P(\eta_k|\eta_j)$. Since $P(\eta_i|\eta_j) = P(\eta_i, \eta_j)/P(\eta_j)$ and $P(\eta_k|\eta_j) = P(\eta_k, \eta_j)/P(\eta_j)$ we get the following approximation for the three-site probability $P(\eta_i, \eta_j, \eta_k) = P(\eta_i, \eta_j)P(\eta_j, \eta_k)/P(\eta_j)$. These results are used in equations (33) and (34) to get a set of closed equations in the variables $P(1) = x$, $P(2) = y$, $P(3) = z$, $P(03) = u$ and $P(12) = v$. With this approximation the model is described by the set of five equations

$$\frac{dx}{dt} = bu - ex, \tag{40}$$

$$\frac{dy}{dt} = -av, \tag{41}$$

$$\frac{dz}{dt} = av - cz, \tag{42}$$

$$\frac{du}{dt} = e(z-u) - (c+rb)u + (1-r)a\frac{(y-v)v}{y} + (1-r)b\frac{u^2}{1-x}, \tag{43}$$

$$\frac{dv}{dt} = -(e+ra)v + (1-r)b\frac{u(y-v)}{1-x} + (1-r)a\frac{v^2}{y}. \tag{44}$$

To obtain the threshold of the spreading of the disease we analyze the stability of the solution $x = 0$, $y = 1$, $z = 1$, $u = 0$ and $v = 0$, which characterizes the state where the disease does not spread. To get the stability of this equation we linearize the evolution around this solution to obtain

$$\frac{dx}{dt} = -ex + bu, \tag{45}$$

$$\frac{d\rho}{dt} = av, \tag{46}$$

$$\frac{dz}{dt} = -cz + av, \tag{47}$$

$$\frac{du}{dt} = ez - (e+c+rb)u + (1-r)av, \tag{48}$$

$$\frac{dv}{dt} = (1-r)bu - (e+ra)v, \tag{49}$$

where $\rho = 1 - y - z$. A linear analysis of stability amounts to calculate the eigenvalues of the matrix composed by the linear coefficients of the right hand side of the above equations,



FIG. 3: Dengue outbreaks according to SMF approximation (a) and PMF approximation (b). In both cases the parameters are $a/c = 10$ and $e/c = 49$ and the initial condition is $x = 10^{-6}$. The figures shows the number of individuals that are being infected per unit time $\zeta = -dy/dt$, or the density of new cases, and the density of infected mosquitoes x as functions of time. The area below the epidemic curve ζ equals the density of recovered individuals ρ_0. For the SMF, $\rho_0 = 0.80$, and for the PMF, $\rho_0 = 0.50$.

given by

$$\begin{pmatrix} -e & 0 & 0 & b & 0 \\ 0 & 0 & 0 & 0 & a \\ 0 & 0 & -c & 0 & a \\ 0 & 0 & e & -(e+c+rb) & (1-r)a \\ 0 & 0 & 0 & (1-r)b & -(e+ra) \end{pmatrix}. \tag{50}$$

Two eigenvalues are $\lambda_1 = 0$ and $\lambda_2 = -e$. The others are the roots of

$$-(\lambda + c)(\lambda + e + c + rb)(\lambda + e + ra) + ea(1-r)b + (\lambda + c)(1-r)^2 ab = 0. \tag{51}$$

The line of stability is obtained by setting in this equation $\lambda = 0$, that is,

$$-c(e + c + rb)(e + ra) + (1-r)abe + (1-r)^2 abc = 0. \tag{52}$$

As before we consider $b = a$ and write the equation that describes the threshold line as

$$(1-r)a^2e + (1-2r)a^2c = e^2c + ec^2 + rac^2 + 2raec. \tag{53}$$

When the coordination number $k = 4$ ($r = 1/4$) the threshold line is described by

$$3a^2e + 2a^2c = 4e^2c + 4ec^2 + ac^2 + 2aec. \tag{54}$$

In figure 2b we show the line described by this equation which represents the phase transition between the spreading to non-spreading regime. Below the transition line the non-spreading solution becomes unstable giving rise to the spreading solution.

The threshold of epidemic obtained by the above linear analysis can equivalently be obtained in terms of the reproductive ratio. In the present case of pair mean-field approach $P(2|1) = P(12)/P(1) = v/x$ and $P(0|3) = P(03)/P(3) = u/z$ so that

$$R_0 = \frac{abuv}{ecxz}. \tag{55}$$

Again, we may say that R_0 depends on the parameters a, b, c and e only through the ratios e/c, b/c and a/c. Indeed, if we substitute λ given by equation (12) into equation (51) we get an equation that gives R_0 in an implicit form. It is easy to see that this equation contains the parameters a, b, c and e only through the ratios e/c, b/c and a/c. Figure 2b shows the lines at constant R_0 in the plane e/c versus a/c for the case $b = a$. We also show the disease threshold line, given by $R_0 = 1$, that separates the non-spreading and the spreading regimes. Comparing the pair and simple mean field approximation we see that the inactive region of the phase diagram is larger for the pair approximation than for the simple approximation as can be seen in figure 2. This aspect can also be observed in figure 3 which shows the epidemic curve for both approximations using the same set of parameters. The simple mean-field predicts a larger number of recovered individuals, and consequently a larger number of individuals that have been infected, as compared to the pair mean-field. As illustrated in figure 3, for the same set of parameters, the final density of recovered individuals are $\rho_0 = 0.80$ for the SMF and $\rho_0 = 0.50$ for the PMF.

We have solved the above set of equations using the initial condition $x = 10^{-6}$, $y = 1$, $z = 0$, $u = 0$ and $v = 10^{-6}$. From the numerical solution we have obtained the epidemic curve, that is the density of individuals that are being infected per unit time, $\zeta = -dy/dt$. Figure 3 shows an example of the epidemic curve obtained in the PMF approximation together with the density of infected mosquitoes.

It is worth mentioning that in the early stages of the spreading of the epidemics the quantities x, z, u and v increase exponentially according to

$$x = x_0 e^{\lambda t}, \qquad z = z_0 e^{\lambda t}, \qquad u = u_0 e^{\lambda t}, \qquad v = v_0 e^{\lambda t}, \tag{56}$$

where λ is the largest eigenvalue of the matrix (50). From this behavior we see that the ratios $v/x = P(2|1)$ and $u/z = P(0|3)$ are independent of time at the early stage of the spreading of disease, and so is the reproductive ratio R_0.

An important aspect of our pair approach concerns the relation of the epidemic curve with the conditional probabilities $P(2|1)$ and $P(0|3)$ which are treated exactly in this approach. Remember that $P(2|1)$ is the conditional probability of the occurrence of a susceptible individual in the presence of a infected mosquito and $P(0|3)$ is the conditional probability of the occurrence of a susceptible mosquito in the presence of a infected individual.

B. Actual epidemics data

The theoretical analysis performed in the previous section was motivated by actual data of two dengue outbreaks, as in a previous work by one of us [10]. They occurred in Salvador, Bahia, Brazil in 1995 and 2002: although in first outbreak no systematic vector population control has been applied by the local health authorities, in the second outbreak a more systematic mosquito control was in course due to the establishment of Brazilian Dengue Control Program. As it was discussed in [10] the transmission force for the second outbreak is more intense than the first outbreak.

The dengue time series of the outbreaks are shown in figure 4. It is easy to observe that the 2002 outbreak (prevalence of DEN3) increases and decreases faster than the 1995 outbreak (only DEN2). It is clear that faster increasing of 2002 outbreak is associated with higher transmission force acquired for DEN3. But here is no consensus whether the faster reduction is more related to transmission force acquired for DEN3 that quickly increased the herd immunity of human population pool, or to vector control actions applied in Salvador in 2002, or both factors [10].

Both time series can be fitted by means of the simple or pair mean-field approximations of our simple model, assuming parameter values for which the active phase is set up. Here we fitted the data from the two outbreaks by means of the pair mean-field approximation, as



FIG. 4: Pair mean-field (PMF) solutions fitted to the dengue outbreaks in Salvador, Brazil, in 1995 (DEN2) and 2002 (DEN3 is prevalent). For both cases we used the same set of parameters, $a = b = 2$ day^{-1}, $e = 9.8$ day^{-1}, $c = 0.2$ day^{-1}. These parameters amounts to a reproductive rate $R_0 = 1.38$.

shown in figure 4. For both outbreaks we used the following values for the three parameters of the model: $a = b = 2$ day^{-1}, $e = 9.8$ day^{-1} and $c = 0.2$ day^{-1}. Using equation (55), these values give a reproductive rate equals $R_0 = 1.38$. The total number of individuals, which can be understood as the carrying capacity, was considered to be 3.2×10^4 for the 1995 outbreak and 6×10^4 for the 2002 outbreak.

V. CONCLUDING REMARKS

In this work we have applied stochastic dynamics to a dengue bipartite lattice model to analyze the transition between epidemic and non-epidemic states in terms of the probability of human-mosquito and mosquito-human transmission and vector control parameters. Analyzing the model by means of simple and pair mean-field approximations for coordination number k, we have shown that a phase transition takes place between epidemic and non-epidemic states for $k \neq 2$. The epidemic curve resulting from the PMF approach were used to fit the the patterns of time series of actual dengue outbreaks that have occurred in Salvador in 1995 and 2002.

We have presented a precise definition of the reproductive rate R_0 which is appropriate for systems described by a stochastic dynamics, that characterizes the spreading of the disease, and we have relate it to the largest eigenvalue of the matrix associated to the evolution equations. This definition can be generalized to other types of disease transmission and seems to be promising in the analysis of epidemics. According to our definition, the reproductive rate is directed related to the conditional probability of the occurrence of a susceptible human (mosquito) given the presence in the neighborhood of an infected mosquito (human). At the early stages of the epidemic, these conditional probabilities are simply equal to the unity in the SMF approach but are nontrivial in the PMF, what makes this approach a richer description when compared to the SMF. Another quantity that characterizes the epidemic is ρ, the quantity that measures the size of the epidemic and vanishes in the nonspreading regime. It was also determined by means of the SMF and PMF approach.

In future work we intend to perform Monte Carlo simulations of the bipartite lattice model in order to analyze the spatial-temporal patterns. They may then be compared with the spatial-temporal patterns of 1995 and 2002 outbreaks which have different features from each other. In another work we will consider a model that takes into account the possibility of the vaccine [24] since there is a perspective of having the vaccine against dengue available within the next five years.

Acknowledgment

STRP, TT and MJO thank CNPq and FAPESB (contract number PNX 006/2009) for supporting the research. SRTP also thanks INCT (contract number 57386/2008-9). TT and MJO acknowledge INCT (contract number 573560/2008-0).

---

[1] D. J. Gubler, "Dengue and dengue hemorrhagic fever: its history and resurgence as a global health problem", in D. J. Gubler and G. Kuno (editors) *Dengue and Dengue hemorrhagic Fever* (CAB International, London, 1997), p. 1-22.

[2] M. J. Keeling and P. Rohani, *Modeling Infectious Diseases in Humans and Animals* (Princeton University Press, Princeton, 2008).

[3] M. J. Keeling, Proc. R. Soc. London B **266**, 859 (1999).

[4] F. Brauer and C. Castillo-Chavez, *Mathematical Models in Population Biology and Epidemiology* (Springer, New York, 2001).

[5] D. R. de Souza and T. Tomé, Physica A **389**, 1142 (2010).

[6] T. Tomé and R. M. Ziff, Phys. Rev. E **82**, 051921 (2010).

[7] T. Tomé and M. J. de Oliveira, J. Phys. A **44**, 095005 (2011).

[8] C. J. Rhodes and R. M. Anderson, Phil. Trans. R. Soc. London B **351**, 1679 (1996).

[9] R. Pastor-Satorras and A. Vespignani, Phys. Rev. Lett. **86**, 3200 (2001).

[10] S. T. R. Pinho, C. P. Ferreira, L. Esteva, F. R. Barreto, V. C. Morato-e-Silva and M. G. L. Teixeira, Phil. Trans. R. Soc. A **368**, 5679 (2010).

[11] N. M. Ferguson, C. A. Donnelly and R. M. Anderson, Phil. Trans. R. Soc. London B **354**, 757 (1999).

[12] E. Massad, F. A. B. Coutinho, M. N. Burattini and L. F. Lopez, Trans. R. Soc. Trop. Med. Hyg. **95**, 370 (2001).

[13] M. L. Martins, G. Ceotto, S. G. Alves, C. C. B. Bufon, J. M. Silva and F. F. Laranjeira, Phys. Rev. E **62**, 7024 (2000).

[14] C. P. Ferreira, P. Pulino, H. M. Yang and L. T. Takahashi, Math. Pop. Studies **13**, 215 (2006).

[15] L. B. L. Santos, M. C. Costa, S. T. R. Pinho, R. F. S. Andrade, F. R. Barreto, M. G. Teixeira and M. L. Barreto, Phys. Rev. E **80**, 016102 (2009).

[16] T. Botari, S. G. Alves and E. D. Leonel, Phys. Rev. E **83**, 037101 (2011).

[17] S. L. Silva, J. A. Ferreira and M. L. Martins, Physica A **377**, 689 (2007).

[18] S. C. Ferreira and M. L. Martins, Phys. Rev. E **76**, 036112 (2007).

[19] F. R. Barreto, M. G. Teixeira, M. D. N. Costa, M. S. Carvalho and M. L. Barreto, BMC Public Health **8**, 51 (2008).

[20] E. A. C. Newton and P. Reiter, Am. J. Trop. Med. Hyg. **47**, 709 (1992).

[21] D. G. Kendall, in M. S. Bartlett, J. R. Stat. Soc. A **120**, 48 (1957).

[22] W. O. Kermack and A. G. McKendrick, Proc. R. Soc. A **115**, 700 (1927).

[23] T. Tomé and K. C. de Carvalho, J. Phys. A **40**, 12901 (2007).

[24] S. S. Whitehead, J. E. Blaney, A. P. Durbin and B. R. Murphy, Nat. Rev. Microbiol. **5**, 518

(2007).